

Class Degree and Relative Maximal Entropy

Mahsa Allahbakhshi and Anthony Quas*

Abstract

Given a factor code π from a one-dimensional shift of finite type X onto an irreducible sofic shift Y , if π is finite-to-one there is an invariant called the degree of π which is defined the number of preimages of a typical point in Y . We generalize the notion of the degree to the class degree which is defined for any factor code on a one-dimensional shift of finite type. Given an ergodic measure ν on Y , we find an invariant upper bound on the number of ergodic measures on X which project to ν and have maximal entropy among all measures in the fibre $\pi^{-1}\{\nu\}$. We show that this bound and the class degree of the code agree when ν is ergodic and fully supported. One of the main ingredients of the proof is a uniform distribution property for ergodic measures of relative maximal entropy.

1 Introduction

It is a well-known result that a 1-dimensional irreducible shift of finite type on a finite alphabet has a unique measure of maximal entropy, the measure developed by Shannon and Parry [15, 20]. In contrast, we consider the relative case in which one is given a factor code $\pi : X \rightarrow Y$ from a shift of finite type X to a sofic shift Y , and a measure ν on Y . In this case ergodic measures on X in the fibre $\pi^{-1}\{\nu\}$ having maximal entropy in the fibre, so-called measures of relative maximal entropy, are not well understood.

Measures of relative maximal entropy appear frequently in different topics in symbolic dynamics, and have been applied to the problem of computing

*We thank the referee for detailed and helpful comments.

the Hausdorff dimension of certain sets by Gatzouras and Peres [7]. There are connections of measures of relative maximal entropy with functions of Markov chains [1, 3, 4, 14], measures that maximize a weighted entropy functional [6, 21], the theory of pressure and equilibrium states [8, 9, 19], relative pressure and relative equilibrium states [11, 12, 23], and compensation functions [3, 23]. Other uses of such measures arise from their application in the mathematics of information transfer [16] and information-compressing channels [14].

Unlike in the case considered by Shannon and Parry where the measure of maximal entropy is unique, it is known that measures of relative maximal entropy may not be unique [16, 17]. It is natural to ask how many of such measures there may be for a particular factor code $\pi : X \rightarrow Y$ and an ergodic invariant measure on Y . A previous result on this problem is the result of Petersen, Quas, and Shin [17] which established an explicit finite upper bound on the number of measures of relative maximal entropy. They considered the case (always achievable by recoding) that π is a 1-block code from a 1-step shift of finite type. In this case, the main result of [17] states that the number of measures of relative maximal entropy is bounded above by the number of preimages of any symbol w of Y which has positive probability with respect to the measure on Y .

The bound in [17] suffers from not being invariant under conjugacy and becomes arbitrarily large simply by recoding to a higher block presentation. To avoid this issue one possibility is to take the minimum of this bound over all shifts of finite type $\tilde{X}$ which are conjugate to X . However, no algorithm for computing this minimum has been found. On the other hand, in this paper we find a simpler conjugacy-invariant upper bound which can be strictly better and is never worse than the minimum-over-conjugates bound mentioned above.

Given a finite-to-one factor code π from a shift of finite type X to a sofic shift Y , the **degree** is defined to be the minimal cardinality of the set of preimages of a point in Y . This is widely-studied and known to be invariant under recoding [13]. One can show that the number of preimages of every transitive point in Y is exactly the degree of π . Previous efforts to find an analogue of this when π is infinite-to-one have proved unsuccessful [2].

Given y in Y , we define an equivalence relation on X which is respected by recoding. The relation is motivated by communicating classes in Markov

chains. Loosely, two points x and x' in $\pi^{-1}(y)$ lie in the same equivalence class “transition class” if one can find an element of $\pi^{-1}(y)$ matching either of x and x' from $-\infty$ to an arbitrarily large positive coordinate n and matching the other at $+\infty$. The minimal number of transition classes (always finite) in $\pi^{-1}(y)$ when y runs over the points in Y is called the **class degree** of the code. We show that the number of transition classes over every transitive point of Y is exactly the class degree. If π is finite-to-one the degree and the class degree of π agree.

We prove an analogue of Theorem 1 in [17]: given an ergodic measure ν on Y , for distinct ergodic measures μ_1 and μ_2 of relative maximal entropy, the relative independent joining of μ_1 and μ_2 assigns measure zero to the set of points (u, v) such that u, v are in a common transition class.

Using a combinatorial characterization of class degree and further arguments, we show that the number of transition classes above a typical point of ν is an upper bound on the number of ergodic measures of maximal entropy in the fibre $\pi^{-1}\{\nu\}$. This bound is equal to the class degree of the code when ν is ergodic and fully supported.

One key ingredient to this work is a relative version of the uniform conditional distribution property for measures of maximal entropy. We follow techniques developed by Burton and Steif [5] (which were related to an earlier work of Lanford and Ruelle [10]) to show a uniform conditional distribution property for measures of relative maximal entropy.

2 Background

Throughout the paper, X will denote a one-dimensional shift of finite type on a finite alphabet (SFT) and the shift map will be denoted by T_X , or simply by T when there is no ambiguity. We will deal with a factor code π from X to a sofic shift Y . A detailed description of these notions is given in [13]. A triple (X, Y, π) is called a **factor triple** when $\pi : X \rightarrow Y$ is a factor code from a SFT X to a sofic shift Y . The alphabet of a shift space X is denoted by $\mathcal{A}(X)$ and the σ -algebra on X generated by cylinder sets of X is denoted by $\mathcal{B}_X$. The set of all n -blocks that occur in points of X is denoted by $\mathcal{B}_n(X)$, and the language of X , $\mathcal{L}(X)$, is the collection $\bigcup_{n=0}^{\infty} \mathcal{B}_n(X)$. Let x be in X and $G \subseteq \mathbb{Z}$. The configuration which occurs in

x on G is denoted by x_G . If $G = [i, j]$ is a $\mathbb{Z}$ -interval we sometimes denote x_G by $x_{[i, j]}$. We say two factor triples (X, Y, π) and $(\tilde{X}, \tilde{Y}, \tilde{\pi})$ are **conjugate**, and denote it by $(X, Y, \pi) \cong (\tilde{X}, \tilde{Y}, \tilde{\pi})$, if X is conjugate to $\tilde{X}$ under a conjugacy ϕ , Y is conjugate to $\tilde{Y}$ under a conjugacy ψ , and $\tilde{\pi} \circ \phi = \psi \circ \pi$. Let (X, Y, π) be a factor triple where π is a 1-block factor code induced by the map $\pi_b : \mathcal{A}(X) \rightarrow \mathcal{A}(Y)$ (b stands for block). The map π_b naturally extends to elements of $\mathcal{L}(X)$. Above every Y -block W of length n there is a set of X -blocks W' of length n which are sent to W by π_b ; i.e., $\pi_b(W') = W$. Given $0 \leq i \leq n - 1$, define

$$d(W, i) = |\{a \in \mathcal{A}(X) : \exists W' \text{ with } \pi_b(W') = W, W'_i = a\}|,$$

and let

$$d_\pi^* = \min\{d(W, i) : W \in \mathcal{L}(Y), 0 \leq i \leq |W| - 1\}.$$

A **magic block** is a block W such that $d(W, i) = d_\pi^*$ for some $0 \leq i \leq |W| - 1$. Such an index i is called a **magic coordinate** of W . A factor code π has a **magic symbol** if there is a magic block of π of length 1.

Theorem 2.1. *[13, Theorem 9.1.7] Let (X, Y, π) be a factor triple. There is a factor triple $(\tilde{X}, \tilde{Y}, \tilde{\pi})$ conjugate to (X, Y, π) such that $\tilde{\pi}$ is a 1-block code with a magic symbol.*

3 Uniform Conditional Distribution

By a well-known result of Parry [15], generalizing an earlier result of Shannon [20], in one dimension every irreducible shift of finite type on a finite alphabet has a unique measure of maximal entropy. Burton and Steif [5] give a counterexample to this statement in higher dimensions. However, they show that such measures all have the uniform conditional distribution property stated in Theorem 3.1 (which is related to an earlier work of Lanford and Ruelle [10]). Given a finite set $G \subseteq \mathbb{Z}^d$, the boundary of the complement of G is $\partial G^c = \{i \in G^c : \exists j \in G \text{ with } \|i - j\| = 1\}$.

Theorem 3.1. *[5, Proposition 1.19] Let X be a 1-step d -dimensional SFT and μ be a measure of maximal entropy on X . Then the conditional distribution of μ on any finite set $G \subseteq \mathbb{Z}^d$ given the configuration on G^c is μ -a.s. uniform over all configurations on G which extend the configuration on ∂G^c .*

Given a factor triple (X, Y, π) and an ergodic measure ν on Y , there can exist more than one ergodic measure of relative maximal entropy over ν ; i.e., there can be more than one ergodic measure on X which projects to ν and has maximal entropy among all measures in the fibre $\pi^{-1}\{\nu\}$, see [17, Example 3.3]. We use Lemma 3.2 and follow techniques developed by Burton and Steif in the proof of Theorem 3.1 to show the uniform conditional distribution property for measures of relative maximal entropy in Theorem 3.3.

Lemma 3.2. [22, Theorem 4.7] *Let $(X, \mathcal{B}, \mu)$ be a probability space. Let $\mathcal{A}$ be a finite sub-algebra of $\mathcal{B}$ and let $(\mathcal{F}_n)_{n=1}^\infty$ be an increasing sequence of sub- σ -algebras of $\mathcal{B}$ with $\bigvee_{n=1}^\infty \mathcal{F}_n = \mathcal{F}$. Then $H(\mathcal{A}|\mathcal{F}_n) \rightarrow H(\mathcal{A}|\mathcal{F})$.*

Theorem 3.3. *Let $\pi : X \rightarrow Y$ be a 1-block factor code from a 1-step, 1-dimensional SFT X to a sofic shift Y . Let ν be an invariant measure on Y , and μ be an invariant measure of relative maximal entropy over ν . Given a finite set $G \subseteq \mathbb{Z}$ and y in Y the conditional distribution of μ on $\pi^{-1}(y)$ restricted to G given the configuration on G^c is μ -a.s. uniform over all configurations on G which extend the configuration on ∂G^c and map to the same configuration in Y under the factor code π .*

Proof. Let G be a finite subset of $\mathbb{Z}$. Let Δ be a configuration of Y on G . Pick a configuration η of X on ∂G^c such that $\mu(\eta \cap \pi^{-1}(\Delta)) > 0$. Starting from μ , we define a measure $\tilde{\gamma}$ on X by uniformizing over pre-images of Δ that have η on the boundary. We then show that if μ does not have the required uniform conditional distribution property then $\tilde{\gamma}$ has greater entropy than μ , but still is an element of the fibre $\pi^{-1}\{\nu\}$. This is a contradiction and will therefore establish the required uniform conditional distribution property of μ .

Let $D = \{\alpha_1, \dots, \alpha_L\}$ be the set of all configurations on G which extend η and map to Δ under the factor code π . Let the $\mathbb{Z}$ -interval $R = [-m, m-1]$ be large enough so that $G \cup \partial G^c \subseteq R$ and let $(R_n)_{n \in \mathbb{Z}}$ be the partition of $\mathbb{Z}$ by translates of R ($R_0 = R$), i.e. $R_n = [(2n-1)m, (2n+1)m-1]$. Let G_n and ∂G_n^c be the corresponding translates of G and ∂G^c in R_n . Given $S \subseteq \mathbb{Z}$ let $\mathcal{P}(X_S)$ be the partition of X generated by the configurations of X on S , and $\sigma(X_S)$ be the σ -algebra generated by $\mathcal{P}(X_S)$. When $S = [a, b]$ is a $\mathbb{Z}$ -interval we sometimes denote X_S by X_a^b . Considering R above, $\sigma(X_{-m}^{m-1})$ is the finite σ -algebra generated by the partition

$$\mathcal{P}(X_{-m}^{m-1}) = \{ -m[x_{-m}x_{-m+1} \dots x_{m-1}]_{m-1} : x_{-m}x_{-m+1} \dots x_{m-1} \in \mathcal{L}(X) \},$$

and $\sigma(X_{-\infty}^{-m-1}) = \sigma(X_{-3m}^{-m-1}) \vee \sigma(X_{-5m}^{-3m-1}) \vee \dots$

We obtain the measure γ from the measures μ and η as follows. Define the map $\Phi : X \times D^{\mathbb{Z}} \rightarrow X$ by

$$\Phi(x, \zeta)_{G_n} = \begin{cases} \zeta_n & \text{if } x_{\partial G_n^c} = \eta \text{ and } x_{G_n} \in D \\ x_{G_n} & \text{otherwise,} \end{cases}$$

and $\Phi(x, \zeta)_{G_n^c \cap R_n} = x_{G_n^c \cap R_n}$ for each integer n . Since ζ_n is in D and each element of D extends η , the assumption that X is a 1-step SFT implies that $\Phi(x, \zeta)$ lies in X . For each ζ in $D^{\mathbb{Z}}$ we have $\pi(x) = \pi(\Phi(x, \zeta))$ since $\Phi(x, \zeta)$ and x are the same except having alternative α_i 's in the same positions. Let C be a cylinder set of X . Define $\gamma(C) = (\mu \times \lambda)\Phi^{-1}(C)$ where λ is the Bernoulli $(1/L, \dots, 1/L)$ measure on $D^{\mathbb{Z}}$. The measure γ is not necessarily invariant under T ; however, for each cylinder set C we have $\gamma(C) = \gamma(T^{-2m}C)$. So the new measure $\tilde{\gamma}$ on X defined by $\tilde{\gamma}(C) = \frac{1}{2m} (\gamma(C) + \dots + \gamma(T^{-2m+1}C))$ is T -invariant. Since for each cylinder set E of Y we have

$$\gamma(\pi^{-1}E) = (\mu \times \lambda)(\pi^{-1}E \times D^{\mathbb{Z}}) = \nu(E),$$

we deduce that both measures γ and $\tilde{\gamma}$ are in the fibre $\pi^{-1}\{\nu\}$.

Define an equivalence relation on X as follows. Given x, x' in X , say $x \sim_0 x'$ if either $x_R = x'_R$ or else $x_{R \cap G^c} = x'_{R \cap G^c}$, $x_{\partial G^c} = x'_{\partial G^c} = \eta$, and x_G, x'_G in D . Denote the equivalence class containing x by $C_0(x)$. Such equivalence classes form a sub-partition of $\mathcal{P}(X_{-m}^{m-1})$. Let $\mathcal{A}$ be the σ -algebra generated by these equivalence classes. We show $h_{\tilde{\gamma}}(T) \geq h_{\mu}(T)$ as follows,

using Lemma 3.4 which appears below.

$$\begin{aligned}
h_{\tilde{\gamma}}(T) &= \frac{1}{2m} h_{\tilde{\gamma}}(T^{2m}) \\
&= \frac{1}{2m} h_{\gamma}(T^{2m}) \\
&= \frac{1}{2m} H_{\gamma}(\sigma(X_{-m}^{m-1}) | \sigma(X_{-\infty}^{-m-1})) \\
&= \frac{1}{2m} H_{\gamma}(\mathcal{A} | \sigma(X_{-\infty}^{-m-1})) + \frac{1}{2m} H_{\gamma}(\sigma(X_{-m}^{m-1}) | \sigma(X_{-\infty}^{-m-1}) \vee \mathcal{A}) \\
&= \frac{1}{2m} H_{\gamma}(\mathcal{A} | \sigma(X_{-\infty}^{-m-1})) + \frac{1}{2m} H_{\gamma}(\sigma(X_{-m}^{m-1}) | \mathcal{A}) && \text{Lem 3.4(a)} \\
&\geq \frac{1}{2m} H_{\gamma}(\mathcal{A} | \sigma(X_{-\infty}^{-m-1})) + \frac{1}{2m} H_{\mu}(\sigma(X_{-m}^{m-1}) | \mathcal{A}) && \text{Lem 3.4(b)} \\
&\geq \frac{1}{2m} H_{\mu}(\mathcal{A} | \sigma(X_{-\infty}^{-m-1})) + \frac{1}{2m} H_{\mu}(\sigma(X_{-m}^{m-1}) | \mathcal{A}) && \text{Lem 3.4(c)} \\
&\geq \frac{1}{2m} H_{\mu}(\mathcal{A} | \sigma(X_{-\infty}^{-m-1})) + \frac{1}{2m} H_{\mu}(\sigma(X_{-m}^{m-1}) | \sigma(X_{-\infty}^{-m-1}) \vee \mathcal{A}) \\
&= \frac{1}{2m} H_{\mu}(\sigma(X_{-m}^{m-1}) | \sigma(X_{-\infty}^{-m-1})) \\
&= \frac{1}{2m} h_{\mu}(T^{2m}) \\
&= h_{\mu}(T).
\end{aligned}$$

Lemma 3.4. *Reusing previous notations, we have*

$$(a) \quad H_{\gamma}(\sigma(X_{-m}^{m-1}) | \sigma(X_{-\infty}^{-m-1}) \vee \mathcal{A}) = H_{\gamma}(\sigma(X_{-m}^{m-1}) | \mathcal{A}).$$

$$(b) \quad H_{\gamma}(\sigma(X_{-m}^{m-1}) | \mathcal{A}) \geq H_{\mu}(\sigma(X_{-m}^{m-1}) | \mathcal{A}). \text{ Equality occurs if and only if}$$

$$\frac{\mu(\alpha \cap \bar{A})}{\mu(\bar{A})} = 1/L,$$

for each α in D and $\bar{A}$ in $\mathcal{P}(\mathcal{A})$ such that for each x in $\bar{A}$ we have $x_{\partial G^c} = \eta$ and x_G in D .

$$(c) \quad H_{\gamma}(\mathcal{A} | \sigma(X_{-\infty}^{-m-1})) \geq H_{\mu}(\mathcal{A} | \sigma(X_{-\infty}^{-m-1})).$$

Proof. By definition, for a measure ρ in $\{\mu, \gamma\}$ we have

$$H_{\rho}(\sigma(X_{-m}^{m-1}) | \mathcal{A}) = - \sum_{i,k} \rho(O_i \cap A_k) \log \frac{\rho(O_i \cap A_k)}{\rho(A_k)}$$

where O_i is in $\mathcal{P}(X_{-m}^{m-1})$ and A_k is in $\mathcal{P}(\mathcal{A})$ (if $\rho(O_i \cap A_k)$ is zero we define $\rho(O_i \cap A_k) \log \frac{\rho(O_i \cap A_k)}{\rho(A_k)} = 0$). Let x be in A_k . If $x_{\partial G^c} \neq \eta$ or $x_G \notin D$ then for every O_i in $\mathcal{P}(X_{-m}^{m-1})$ we have either $O_i \cap A_k = \emptyset$ or $O_i \cap A_k = A_k$ which both imply $\rho(O_i \cap A_k) \log \frac{\rho(O_i \cap A_k)}{\rho(A_k)} = 0$. Let $\{\bar{A}_1, \dots, \bar{A}_M\}$ be the set consisting of elements in the partition $\mathcal{P}(\mathcal{A})$ so that for any point x in $\bar{A}_k$ we have $x_{\partial G^c} = \eta$ and x_G in D . Then

$$H_\rho(\sigma(X_{-m}^{m-1})|\mathcal{A}) = - \sum_{i,k} \rho(O_i \cap \bar{A}_k) \log \frac{\rho(O_i \cap \bar{A}_k)}{\rho(\bar{A}_k)}.$$

There are exactly L pairwise disjoint sets O_i in $\mathcal{P}(X_{-m}^{m-1})$ defined by blocks which agree everywhere except on G , and form a partition of $\bar{A}_k$. Let these sets be denoted by $O_{k,1}, \dots, O_{k,L}$ where $O_{k,i} = \alpha_i \cap \bar{A}_k$ for each $1 \leq i \leq L$. It follows that

$$\begin{aligned} H_\rho(\sigma(X_{-m}^{m-1})|\mathcal{A}) &= - \sum_{k,i} \rho(O_{k,i} \cap \bar{A}_k) \log \frac{\rho(O_{k,i} \cap \bar{A}_k)}{\rho(\bar{A}_k)} \\ &= - \sum_{k,i} \rho(\alpha_i \cap \bar{A}_k) \log \frac{\rho(\alpha_i \cap \bar{A}_k)}{\rho(\bar{A}_k)}. \end{aligned} \tag{3.2}$$

Let $n \geq 1$. By definition of γ , for each configuration α in D , each cylinder set P in $\mathcal{P}(X_{-(2n+1)m}^{-m-1})$, and each $1 \leq k \leq M$ we have

$$\frac{\gamma(\alpha \cap P \cap \bar{A}_k)}{\gamma(P \cap \bar{A}_k)} = \frac{\gamma(\alpha \cap \bar{A}_k)}{\gamma(\bar{A}_k)} = \frac{1}{L}.$$

It follows that

$$\begin{aligned} H_\gamma(\sigma(X_{-m}^{m-1})|\sigma(X_{-(2n+1)m}^{-m-1}) \vee \mathcal{A}) &= - \sum_{i,j,k} \gamma(\alpha_i \cap P_j \cap \bar{A}_k) \log \frac{\gamma(\alpha_i \cap P_j \cap \bar{A}_k)}{\gamma(P_j \cap \bar{A}_k)} \\ &= \log L \sum_{i,k} \gamma(\alpha_i \cap \bar{A}_k) \\ &= H_\gamma(\sigma(X_{-m}^{m-1})|\mathcal{A}) \end{aligned}$$

(use the same argument we had before Equation (3.2) to get the first equality above). Then (a) follows from Lemma 3.2 and the fact that $\left(\sigma(X_{-(2n+1)m}^{-m-1})\right)_{n=1}^\infty$ is an increasing sequence of σ -algebras with $\bigvee_{n=1}^\infty \sigma(X_{-(2n+1)m}^{-m-1}) = \sigma(X_{-\infty}^{-m-1})$.

(b) is a standard property of entropy.

To show (c) let $\mathcal{A}_{-(2n+1)m}^{-m-1}$ denote the σ -algebra $\bigvee_{i=1}^n T^{-2mi} \mathcal{A}$ (the choice of notation is intended to remind the reader that $\mathcal{A}_{-\infty}^{-m-1}$ is a sub- σ -algebra of $\sigma(X_{-\infty}^{-m-1})$).

Any set B in $\mathcal{A}_{-(2n+1)m}^{-m-1}$ splits up into L^K elements of $\sigma(X_{-(2n+1)m}^{-m-1})$, $B_1, \dots, B_{L^K}$ of equal measure (where K is the number of occurrences in B of blocks that are randomized under γ). Each of these sets has the property that $\gamma(A \cap B_i)/\gamma(B_i) = \gamma(A \cap B)/\gamma(B)$. Accordingly we see that $H_\gamma(\mathcal{A}|\mathcal{A}_{-(2n+1)m}^{-m-1}) = H_\gamma(\mathcal{A}|\sigma(X_{-(2n+1)m}^{-m-1}))$. Taking a limit we obtain

$$H_\gamma(\mathcal{A}|\sigma(X_{-\infty}^{-m-1})) = H_\gamma(\mathcal{A}|\mathcal{A}_{-\infty}^{-m-1}).$$

We then notice that $H_\gamma(\mathcal{A}|\mathcal{A}_{-\infty}^{-m-1}) = H_\mu(\mathcal{A}|\mathcal{A}_{-\infty}^{-m-1})$ since the measures μ and γ agree on elements of $\bigvee_{i=0}^\infty T^{-2mi} \mathcal{A}$.

Finally it is clear that $H_\mu(\mathcal{A}|\mathcal{A}_{-\infty}^{-m-1}) \geq H_\mu(\mathcal{A}|\sigma(X_{-\infty}^{-m-1}))$. Combining these three equalities and inequalities we obtain $H_\gamma(\mathcal{A}|\sigma(X_{-\infty}^{-m-1})) \geq H_\mu(\mathcal{A}|\sigma(X_{-\infty}^{-m-1}))$ as required. $\square$

Proof of Theorem 3.3 (continued). Since μ is a measure of relative maximal entropy it follows that all of the inequalities in Equations (3.1) are forced to be equalities. In particular, we have $H_\mu(\sigma(X_{-m}^{m-1})|\mathcal{A}) = H_\gamma(\sigma(X_{-m}^{m-1})|\mathcal{A})$. Then Lemma 3.4(b) implies that

$$\frac{\mu(\alpha \cap \bar{A})}{\mu(\bar{A})} = 1/L, \quad (3.3)$$

for each configuration α in D and $\bar{A}$ in $\mathcal{P}(\mathcal{A})$ such that for each x in $\bar{A}$ we have $x_{\partial G^c} = \eta$ and x_G in D . Note that given a finite set $G \subseteq \mathbb{Z}$, both configurations $\Delta \in \mathcal{B}_Y$ on G and $\eta \in \mathcal{B}_X$ on ∂G^c with $\mu(\eta \cap \pi^{-1}(\Delta)) > 0$ are chosen arbitrarily. By choosing different configurations and noting that $\bar{A} \in \mathcal{P}(\pi^{-1}(\sigma(Y_G)) \vee \sigma(X_{R \cap G^c})) = \mathcal{P}(\pi^{-1}(\sigma(Y_R)) \vee \sigma(X_{R \cap G^c}))$, Equation (3.3) implies that for any configuration α of X occurring at G we have

$$\mathbb{E}(1_\alpha | \pi^{-1}(\sigma(Y_R)) \vee \sigma(X_{R \cap G^c}))(x) = \begin{cases} \frac{1}{L_{(x,G)}} & \alpha \text{ extends } x_{\partial G^c}, x_G \in \pi_b^{-1}(\pi_b(\alpha)) \\ 0 & \text{otherwise.} \end{cases}$$

where $L_{(x,G)}$ is the number of configurations of X occurring at G which extend $x_{\partial G^c}$ and project to $\pi_b(x_G)$.

Now for a positive integer t let $R^{(t)} = [-(m+t), m+t-1]$, and let $(R_n^{(t)})_{n \in \mathbb{Z}}$ be the partition of $\mathbb{Z}$ by translates of $R^{(t)}$ where $R_0^{(t)} = R^{(t)}$, i.e. $R_n^{(t)} = [(2n-1)(m+t), (2n+1)(m+t)-1]$. Let $R^{(0)} = R$. Since $(\sigma(Y_{R^{(t)}}))_{t=0}^\infty$ and $(\sigma(X_{R^{(t)} \cap G^c}))_{t=0}^\infty$ are increasing sequences with $\bigvee_{t=0}^\infty \sigma(Y_{R^{(t)}}) = \mathcal{B}_Y$ and $\bigvee_{t=0}^\infty \sigma(X_{R^{(t)} \cap G^c}) = \sigma(X_{G^c})$ it follows from Lemma 3.2 that

$$\mathbb{E}(1_\alpha | \pi^{-1}(\mathcal{B}_Y) \vee \sigma(X_{G^c}))(x) = \begin{cases} \frac{1}{L(x, G)} & \alpha \text{ extends } x_{\partial G^c}, x_G \in \pi_b^{-1}(\pi_b(\alpha)) \\ 0 & \text{otherwise.} \end{cases}$$

□

4 Class Degree of a factor code

When π is a finite-to-one factor code from a SFT X to a sofic shift Y , there is a uniform upper bound on the number of pre-images of points in Y [13]. The minimal number of π -pre-images of points in Y is called the **degree** of the code and denoted by d_π .

Theorem 4.1. [13] *Let $\pi : X \rightarrow Y$ be a finite-to-one factor code from a SFT X to an irreducible sofic shift Y . Then every doubly transitive point of Y has exactly d_π preimages.*

Theorem 9.1.11 in [13] shows this result when X is irreducible. However, the proof only depends on the irreducibility of Y .

Corollary 4.2. *Let (X, Y, π) and $(\tilde{X}, \tilde{Y}, \tilde{\pi})$ be conjugate factor triples. Then we have $d_\pi = d_{\tilde{\pi}}$.*

Definition 4.3. *A family $x^{(1)}, \dots, x^{(k)}$ of sequences in a SFT X is **mutually separated** if, for each integer i , $x_i^{(1)}, \dots, x_i^{(k)}$ are all distinct.*

Proposition 4.4. [13] *Let $\pi : X \rightarrow Y$ be a finite-to-one 1-block factor code from a SFT to an irreducible sofic shift Y . The preimages of a transitive point are mutually separated. If π has a magic symbol w then $d_\pi = |\pi_b^{-1}(w)|$.*

For a proof, the reader is referred to Proposition 9.1.9 and Exercise 9.1.3 of [13].

We will find a quantity analogous to the degree when π is an infinite-to-one factor code. This will be done by developing the following equivalence relation on X . Figure 1 illustrates Definition 4.5.

Definition 4.5. Suppose (X, Y, π) is a factor triple and $x, x' \in X$. We say there is a **transition** from x to x' and denote it by $x \rightarrow x'$ if for each integer n , there exists a point v in X so that

1. $\pi(v) = \pi(x) = \pi(x')$, and
2. $v_{-\infty}^n = x_{-\infty}^n$, $v_i^\infty = x_i'^\infty$ for some $i \geq n$.

Write $x \nrightarrow x'$ if the above conditions do not hold. We write $x \sim x'$, and say x and x' are in the same **transition class** if $x \rightarrow x'$ and $x' \rightarrow x$. The relation $\sim$ is an equivalence relation. Denote the set of transition classes in X over $y \in Y$ by $\mathcal{C}_\pi(y)$. Sometimes we denote $\mathcal{C}_\pi(y)$ simply by $\mathcal{C}(y)$ when there is no ambiguity in understanding π . Say $[x] \rightarrow [x']$ if $x \rightarrow x'$ (well-defined). Use the notation $[x] \nrightarrow [x']$ otherwise.




Figure 1: Transition from point x to x'

The following fact is derived from Definition 4.5 immediately.

Fact 4.6. Let $\pi : X \rightarrow Y$ be a 1-block factor code from a 1-step SFT X to a sofic shift Y . Let x, x' project to the same point under π and $x_{a_i} = x'_{a_i}$ where $(a_i)_{i \in \mathbb{N}}$ is a strictly increasing sequence in $\mathbb{Z}$. Then we have $x \sim x'$.

Note that Fact 4.6 gives an obvious case when two points lie in the same transition class. Simple examples allow one to find equivalent points without having a common symbol at the same time.

Definition 4.7. Let (X, Y, π) be a factor triple. The minimal number of transition classes over points of Y is called the **class degree** of π and denoted by c_π .

The following proposition shows that conjugate factor triples have the same class degree.

Proposition 4.8. *Let (X, Y, π) and $(\tilde{X}, \tilde{Y}, \tilde{\pi})$ be conjugate factor triples under conjugacies $\phi : X \rightarrow \tilde{X}$ and $\psi : Y \rightarrow \tilde{Y}$. Then*

- (a) *For each y in Y , $|\mathcal{C}_\pi(y)| = |\mathcal{C}_{\tilde{\pi}}(\psi(y))|$.*
- (b) *$c_\pi = c_{\tilde{\pi}}$.*

Proof. (a) is clear by noting that given u, v in X , $\phi(u) \sim \phi(v)$ if and only if $u \sim v$, and (b) is a direct corollary of (a). $\square$

The following two theorems give an upper bound on the number of transition classes over points of Y . Later in Theorem 4.22 we give a finitary characterization of the number of such classes.

Theorem 4.9. *Let $\pi : X \rightarrow Y$ be a 1-block factor code from a 1-step SFT X to a sofic shift Y . Let y in Y and let w be a symbol in Y occurring infinitely often in positive coordinates of y . Then $|\mathcal{C}(y)| \leq |\pi_b^{-1}(w)|$.*

Proof. Let $(a_j)_{j \in \mathbb{N}}$ be a strictly increasing sequence of integers such that $y_{a_j} = w$ for each $j \in \mathbb{N}$. Let $|\pi_b^{-1}(w)| = d$ and suppose $\pi^{-1}(y)$ contains $d + 1$ distinct transition classes $C_1, \dots, C_{d+1}$. Form a set $A = \{x_1, \dots, x_{d+1}\}$ where x_i is an arbitrary point of C_i . Since $\pi_b^{-1}(w)$ contains exactly d symbols, it follows that for each $j \in \mathbb{N}$ there are at least two points in A with the same a_j th coordinate. The Pigeonhole Principle implies that there is a subsequence $(b_k)_{k \in \mathbb{N}}$ of $(a_j)_{j \in \mathbb{N}}$ and at least two points x and x' in A with $x_{b_k} = x'_{b_k}$ for each $k \in \mathbb{N}$. Fact 4.6 implies that $x \sim x'$. This contradicts the assumption that x and x' are in different transition classes. $\square$

A similar proof to the above implies the following theorem.

Theorem 4.10. *Let $\pi : X \rightarrow Y$ be a 1-block factor code from a 1-step SFT X to a sofic shift Y . Then $|\mathcal{C}(y)| < \infty$ for each $y \in Y$. Moreover, $\min_{w \in \mathcal{A}(Y)} \{|\pi_b^{-1}(w)|\}$ is the uniform upper bound on the number transition classes over each right transitive point of Y .*

We show when $\pi : X \rightarrow Y$ is a factor code from a SFT X to an irreducible sofic shift Y , there are exactly c_π transition classes over a transitive point of Y . In order to show this, we introduce another quantity c_π^* in Definition 4.12, defined concretely in terms of blocks. Since the class degree is invariant under conjugacy of factor triples, by recoding we may focus only on 1-step SFTs and 1-block factor codes.

Definition 4.11. Let $\pi : X \rightarrow Y$ be a 1-block factor code from a 1-step SFT X to a sofic shift Y . Let W be a Y -block of length $p + 1$. Let n be an integer in $(0, p)$, and M be a subset of $\pi_b^{-1}(W_n)$. We say $U \in \pi_b^{-1}(W)$ is *routable through* $a \in M$ *at time* n if there is a block $U' \in \pi_b^{-1}(W)$ with $U'_0 = U_0$, $U'_n = a$, and $U'_p = U_p$. A triple (W, n, M) is called a **transition block** of π if every block in $\pi_b^{-1}(W)$ is routable through a symbol of M at time n . The cardinality of the set M is called the **depth** of the transition block (W, n, M) .

Figure 2 illustrates Definition 4.11.




Figure 2: (W, n, M) is a transition block with $M = \{a_1, a_2\}$. The blocks $U, V, K \in \pi_b^{-1}(W)$ are routable through members of M at time n via blocks $U', V', K' \in \pi_b^{-1}(W)$.

Definition 4.12. Let

$$c_\pi^* = \min\{|M| : (W, n, M) \text{ is a transition block of } \pi\}.$$

A *minimal transition block* of π is a transition block of depth c_π^* .

Example 4.13. In figure 3, we display an example of a labeled graph which defines an infinite-to-one 1-block factor code π . We see that $(001, 1, \{b\})$ is a minimal transition block of π of depth 1. For example, observe that block $U = aac$ is routable through b at time 1 by considering $U' = abc$.




Figure 3: Graph for Example 4.13

We need to develop some definitions and lemmas before proving Theorem 4.22 below. Given a factor triple (X, Y, π) , a point y in Y and a block V occurring in y , it may be the case that some preimages of V can not be extended to a point in X which projects to y under π . Here we show that in order to recognize those non-extendable blocks we do not need to deal with a bi-infinite sequence. A finite extension of a block is sufficient to determine if the block can be extended to a point above y .

Definition 4.14. Let (X, Y, π) be a factor triple and y be in Y . Let V be a block of Y occurring in y at a $\mathbb{Z}$ -interval $[m, n]$. A Block W of Y is called a *y-synchronizing extension* of V if W occurs in y at $[m - l, n + l]$ for some integer $l \geq 0$ and

$$\begin{aligned} \{U \in \pi_b^{-1}(V) : \exists x \in \pi^{-1}(y), x_{[m, n]} = U\} \\ = \{U \in \pi_b^{-1}(V) : \exists B \in \pi_b^{-1}(W), B_{[l, l+n-m]} = U\}. \end{aligned}$$

Lemma 4.15. Let (X, Y, π) be a factor triple and y in Y . Let V be a block occurring in y at a $\mathbb{Z}$ -interval $[m, n]$. There is a *y-synchronizing extension* of V .

Proof. Let

$$S = \{U \in \pi_b^{-1}(V) : \exists x \in \pi^{-1}(y), x_{[m, n]} = U\}.$$

For $l \geq 0$ let

$$S^l = \{U \in \pi_b^{-1}(V) : \exists B^l \in \pi_b^{-1}(y_{[m-l, n+l]}), B^l_{[l, l+n-m]} = U\}.$$

We show for some l we have $S = S^l$. Then it will imply that $y_{[m-l, n+l]}$ is a *y-synchronizing extension* of V .

Clearly for all $l \geq 0$ we have $S \subseteq S^l$. We show $S^l \subseteq S$ for some l as follows. First note that since π is a factor code, S is not empty. It follows that the sequence $(S^l)_{l=0}^\infty$ is a non-increasing sequence of non-empty finite sets. Thus there is l such that for each $r \geq l$ we have $S^l = S^r$. We show $S^l \subseteq S$.

Let U be in S^l . Then U is in S^r for each $r \geq l$. It follows that for each $r \geq l$ there is B^r in $\pi_b^{-1}(y_{[m-r, n+r]})$ with $B_{[r, r+n-m]}^r = U$. For each $r \geq l$ consider the cylinder set $C_{m-r}(B_r)$. The sequence $(C_{m-r}(B_r))_{r \geq l}$ is a non-increasing sequence of non-empty compact subsets of X . It follows that $\bigcap_{r \geq l} C_{m-r}(B_r)$ is not empty. Let $x \in \bigcap_{r \geq l} C_{m-r}(B_r)$. Then $x_{[m, n]} = U$ and $\pi(x) = y$ which implies that U is in S . $\square$

Here we introduce a relation between symbols and transition classes.

Definition 4.16. Let $\pi : X \rightarrow Y$ be a 1-block factor code from a 1-step SFT X to a sofic shift Y . Let y be in Y . Let n be in $\mathbb{Z}$, i be a symbol in $\pi_b^{-1}(y_n)$, and C be a transition class in $\mathcal{C}(y)$. Say i **belongs to C at time n** , and denote it by $i \in S_n(C)$, if

$$\{D \in \mathcal{C}(y) : \text{there is } x \text{ in } D \text{ with } x_n = i\} = \{D \in \mathcal{C}(y) : C \rightarrow D\}.$$

In other words, the set of transition classes in $\mathcal{C}(y)$ containing a point with n th coordinate being i , is the same as the set of transition classes to which there is a transition from the class C (including C itself).

Say i is **transient at time n** if there is no transition class C in $\mathcal{C}(y)$ for which $i \in S_n(C)$.

Example 4.17 illustrates the definition above.

Example 4.17. Consider the directed labeled graph in Figure 4 which presents a 1-block factor code $\pi : X \rightarrow Y$ where $X \subseteq \{a, b, c, d, e, f, g\}^\mathbb{Z}$ and $Y \subseteq \{0, 1\}^\mathbb{Z}$. Let y be the point $\cdots 0101010 \cdots$ in Y . By Definition 4.5, there are 3 distinct transition classes in $\mathcal{C}(y)$ as follows:

1. Transition Class $C_1 = [x_1] = \{x_1\}$ where $x_1 = \cdots bababab \cdots$.
2. Transition Class $C_2 = [x_2]$ where x_2 is a point in $\pi^{-1}(y)$ which does not contain symbols e or g but contains only d 's and f 's from some time onwards, for example: $x_2 = \cdots babcdf d \cdots$.




Figure 4: Graph for Example 4.17

3. *Transition Class $C_3 = [x_3]$ where x_3 is a point in $\pi^{-1}(y)$ which does not contain symbols d or f but contains only e 's and g 's from some time onwards, for example: $x_3 = \cdots egegege \cdots$.*

Clearly $C_1 \rightarrow C_2$ and $C_1 \rightarrow C_3$, but not vice versa. $C_2 \nrightarrow C_3$, $C_3 \nrightarrow C_2$, and symbol c is transient at any time.

Note that the class degree of the factor code π is 2 (consider a point of Y in which the block 11 occurs at some time).

Lemma 4.18. *Let $\pi : X \rightarrow Y$ be a 1-block factor code from a 1-step SFT X to a sofic shift Y , and x be a point in X . There is $m < \infty$ such that for each $n \geq m$ the symbol x_n is not transient. In fact, m can be found in such a way that for each $n \geq m$, x_n belongs to the class $[x]$ at time n ; i.e., $x_n \in S_n[x]$.*

Proof. Let $\pi(x) = y$. Suppose that the transition class C in $\mathcal{C}(y)$ satisfies $[x] \nrightarrow C$. Then there exists $i < \infty$ such that for any z in $\pi^{-1}(y)$ with $z_i = x_i$ we have $z \notin C$. Denote the smallest such i by i^C . Note that i^C may be $-\infty$ (but not $+\infty$). Let $m = \max\{i^C : C \in \mathcal{C}(y), [x] \nrightarrow C\}$. Let $n \geq m$. We show

$$\{C \in \mathcal{C}(y) : [x] \rightarrow C\} = \{C \in \mathcal{C}(y) : \text{there is } z \in C \text{ with } z_n = x_n\}, \quad (4.1)$$

which will imply the result $x_n \in S_n[x]$.

Let C be a transition class in $\mathcal{C}(y)$ with $[x] \rightarrow C$. It follows that $x \rightarrow z$ for some z in C . Therefore, there is a point u in $\pi^{-1}(y)$ as follows.

$$u_t = \begin{cases} x_t & \text{if } -\infty < t \leq n \\ z_t & \text{if } l \leq t < \infty, \end{cases}$$

for some $l \geq n$. Since $u_t = z_t$ from time l onwards, it follows that u is in the class C . Moreover, we have $u_n = x_n$ by the construction. It follows that C belongs to the set in the right hand side of Equation (4.1).

Now let C be an equivalence class in $\mathcal{C}(y)$ with $[x] \not\rightarrow C$. Suppose, for a contradiction, C contains a point z with $z_n = x_n$. Form the point u as follows.

$$u_t = \begin{cases} x_t & \text{if } -\infty < t \leq n \\ z_t & \text{if } n < t < \infty. \end{cases}$$

Note that since X is a 1-step SFT and $z_n = x_n$, the point u is in X . Moreover u maps to y by the construction. Noticing that $u_t = z_t$ from the time n onwards, shows that u lies in the class C . Recall that $n \geq i^C$. By the argument in the first paragraph of the proof, any point in $\pi^{-1}(y)$ with i^C th coordinate equal to x_{i^C} does not belong to C . This contradicts that u is a point in C . $\square$

Lemma 4.19. *Let $\pi : X \rightarrow Y$ be a 1-block factor code from a 1-step SFT X to a sofic shift Y . Let $y \in Y$. There is $m < \infty$ such that for each $n \geq m$ and C in $\mathcal{C}(y)$ there is a symbol i in $\pi_b^{-1}(y_n)$ which belongs to the class C at time n ; i.e., $i \in S_n(C)$.*

Proof. Let $\mathcal{C}(y) = \{C_1, \dots, C_d\}$ for some finite d . Let $A = \{x^{(1)}, \dots, x^{(d)}\}$ be a set containing an arbitrary point $x^{(i)}$ of C_i for each C_i . By Lemma 4.18 there is a $n_{C_i} < \infty$ such that for each $n \geq n_{C_i}$ the symbol $x_n^{(i)}$ belongs to C_i at time n . Let $m = \sup\{n_{C_i} : 1 \leq i \leq d\}$. Note that $m < \infty$. Moreover, for each $n \geq m$ and C_i the symbol $x_n^{(i)}$ belongs to the class C_i at time n . $\square$

Proposition 4.20. *Let $\pi : X \rightarrow Y$ be a 1-block factor code from a 1-step SFT X to a sofic shift Y . Let ν be an invariant measure on Y . Then*

$$\nu(\{y \in Y : \forall C \in \mathcal{C}(y), \forall n \in \mathbb{Z} \text{ there is } i \in \pi_b^{-1}(y_n) \text{ with } i \in S_n(C)\}) = 1.$$

Proof. Let y in Y and let $m(y)$ be the infimum of the set of m 's with the properties given in the statement of Lemma 4.19. We show

$$\nu(\{y \in Y : m(y) = -\infty\}) = 1$$

which will imply the result immediately. Note that $C \in \mathcal{C}(y)$ if and only if $T(C) \in \mathcal{C}(T(y))$. This shows $m(T(y)) = m(y) - 1$. For $k < \infty$ let $A_k = \{y \in Y : m(y) = k\}$. We have $T(A_k) = A_{k-1}$. Since ν is T -invariant it follows that $\nu(A_k) = 0$. Therefore $m(y) = -\infty$ for ν -a.e. y in Y . $\square$

Definition 4.21. Let X be a shift space. A point x in X is **recurrent** if for each block B which occurs in x , B occurs infinitely often in the positive coordinates of x .

Note that given any invariant measure μ on X , μ -almost every point of X is recurrent.

Theorem 4.22. Let π be a 1-block factor code from a 1-step SFT X to a sofic shift Y . Let y be a point of Y and d be the minimal depth of a transition block occurring in y . Then $d \leq |\mathcal{C}(y)|$. Moreover, if y is recurrent then $d = |\mathcal{C}(y)|$.

Proof. We show $d \leq |\mathcal{C}(y)|$ by finding a transition block of depth $|\mathcal{C}(y)|$ occurring in y . We construct such a transition block in the following 4 stages. Let $h = |\mathcal{C}(y)|$ and suppose $\mathcal{C}(y) = \{C_1, \dots, C_h\}$. Choose an integer n_1 satisfying the properties given in the statement of Lemma 4.19.

Stage 1. We claim there is $n_2 \in [n_1, \infty)$ such that for each $x \in \pi^{-1}(y)$, there is $n_1 \leq t \leq n_2$ so that x_t is not transient.

Proof. Suppose there is no such n_2 . It follows that for each $j \geq n_1$ there is a point $x^{(j)}$ in $\pi^{-1}(y)$ such that $x_l^{(j)}$ is transient for all $n_1 \leq l \leq j$. Consider the sequence $(x^{(j)})_{j \in \mathbb{Z}}$, and let x be the limit of a convergent subsequence of it. Clearly $x \in \pi^{-1}(y)$. However, x_l is transient for each $l \geq n_1$, contradicting Lemma 4.18. $\square$

Stage 2. We claim there is $n_3 \in [n_2, \infty)$ and a set of symbols

$$M' = \{a_1, \dots, a_h\}$$

with $a_e \in S_{n_3}(C_e)$ such that for each i in $S_{n_2}(C_e)$ there is a block

$$U \in \pi_b^{-1}(y_{n_2}y_{n_2+1} \dots y_{n_3-1}y_{n_3})$$

which begins with i and ends with a_e . See Figure 5.

Proof. Let C_e be a transition class in $\mathcal{C}(y)$ and x^e be a point in C_e . For each symbol i in $S_{n_2}(C_e)$ (non-empty by the choice of n_1), there is a bi-infinite sequence z^i in the class C_e with $z_{n_2}^i = i$ such that z^i matches x^e from some time $k^i \in [n_2, \infty)$ onwards. Let

$$k^e = \max\{k^i : i \in S_{n_2}(C_e)\},$$




Figure 5: An example illustrating Stage 2. $\mathcal{C}(y) = \{C_1, C_2, C_3\}$, $i_1, i_2, i_3 \in S_{n_2}(C_1)$, $i_4 \in S_{n_2}(C_2)$, $i_5, i_6 \in S_{n_2}(C_3)$, and $M' = \{a_1, a_2, a_3\}$.

and

$$n_3 = \max\{k^e : C_e \in \mathcal{C}(y)\}.$$

Note that $n_3 < \infty$. Rename $x_{n_3}^e$ for each $C_e \in \mathcal{C}(y)$ as a_e , and let $M' = \{a_1, \dots, a_h\}$. $\square$

Stage 3. We claim there is $n_4 \in [n_3, \infty)$ such that for each point x in $\pi^{-1}(y)$ there is a point x' in $\pi^{-1}(y)$ so that $x'_r = x_r$ for every $r \in (-\infty, n_1] \cup [n_4, \infty)$, and $x'_{n_3} \in M'$. See Figure 6.

Proof. Let x be a point in $\pi^{-1}(y)$. By stage 1, there is $n_1 \leq t \leq n_2$ such that x_t is not transient; i.e., x_t belongs to some transition class C_e at time t . It follows that there is a point u in the class C_e with $u_t = x_t$. The new block

$$\dots x_{n_1} \dots x_t u_{t+1} \dots u_{n_2} \tag{4.2}$$

is an allowable block of X . Moreover, this block maps to the block

$$\dots y_t \dots y_{n_2}.$$

Note that the symbol u_{n_2} belongs to the class C_e at time n_2 . So by Stage 2, there is a block starting at u_{n_2} ending at a_e which maps to

$$y_{n_2} \dots y_{n_3}.$$




Figure 6: Graph for Stage 3. $M' = \{a_1, a_2, a_3\}$. $x, x' \in \pi^{-1}(y)$, $x'_r = x_r$ for each $r \in (-\infty, n_1] \cup [n_4, \infty)$, and $x'_{n_3} \in M'$. Same for z, z' and u, u' .

Denote this block by

$$u_{n_2}v_{n_2+1} \dots v_{n_3-1}a_e. \quad (4.3)$$

Connect blocks in Equations (4.2) and (4.3) at u_{n_2} to get the following allowable block of X

$$\dots x_{n_1} \dots x_t u_{t+1} \dots u_{n_2} v_{n_2+1} \dots v_{n_3-1} a_e$$

Observe that having x_t in $S_t(C_e)$ implies that x must belong to a transition class C_f with $C_e \rightarrow C_f$. On the other hand having a_e in $S_{n_3}(C_e)$ implies that there is a point b in C_e with

$$b_{(-\infty, n_3]} = \dots x_{n_1} \dots x_t u_{t+1} \dots u_{n_2} v_{n_2+1} \dots v_{n_3-1} a_e$$

which matches x from some time j onwards for some $n_3 \leq j < \infty$. Let

$$n_x = \min_{n_3 \leq j < \infty} \{j : \exists x' \in \pi^{-1}(y) \text{ with } x'_r = x_r \forall r \in (-\infty, n_1] \cup [j, \infty), \ x'_{n_3} = a_e\}.$$

We claim that there is $n^e < \infty$ such that for each $x \in \pi^{-1}(y)$ with $x_t \in S_t(C_e)$ for some $n_1 \leq t \leq n_2$, there exists a point $x' \in \pi^{-1}(y)$ with $x'_r = x_r$ for each $r \in (-\infty, n_1] \cup [n^e, \infty)$, and $x'_{n_3} = a_e$. Then letting

$$n_4 = \max\{n^e : C_e \in \mathcal{C}(y)\} < \infty$$

will complete the proof of Stage 3.

Suppose, for a contradiction, that there does not exist such an n^e . Then there is a sequence $x^{(l)}$ with $x_{t_l}^{(l)} \in S_{t_l}(C_e)$ for some $n_1 \leq t_l \leq n_2$, such that $\lim_{l \rightarrow \infty} n_{x^{(l)}} = \infty$. Let x^* be the limit of a convergent subsequence of $x^{(l)}$. Clearly x^* is in $\pi^{-1}(y)$ and $n_{x^*} = \infty$ which contradicts the fact that for each x in $\pi^{-1}(y)$, $n_x < \infty$. $\square$

Stage 4. Let W be a y -synchronizing extension of the block

$$V = y_{n_1} \cdots y_{n_4}$$

occurring in $[n_0, n_5]$ for some $n_0 \leq n_1$ and $n_5 \geq n_4$. We claim $(W, n_3 - n_0, M')$ is a transition block of depth $|\mathcal{C}(y)|$.

Proof. Let $U \in \pi_b^{-1}(W)$. By the definition of y -synchronizing extension, there is x in $\pi^{-1}(y)$ with $x_{[n_1, n_4]} = U$. By Stages 1, 2, and 3 there is a point x' in $\pi^{-1}(y)$ with $x'_r = x_r$ for each $r \in (-\infty, n_1] \cup [n_4, \infty)$, and $x'_{n_3} \in M'$. Having $x'_{n_1} = x_{n_1}$, $x'_{n_3} \in M'$, $x'_{n_4} = x_{n_4}$ and $\pi_b(x'_{[n_1, n_4]}) = V$ simply means that U is routable through a symbol of M' at time $n_3 - n_0$. Since U in $\pi_b^{-1}(W)$ is arbitrary it follows that $(W, n_3 - n_0, M')$ is a transition block. Because M' has $|\mathcal{C}(y)|$ elements it follows that W is a transition block of depth $|\mathcal{C}(y)|$. $\square$

Now suppose y is recurrent. Then a transition block of depth d which occurs in y , it occurs infinitely often in the positive coordinates of y . Apply the same method which was used in the proof of Theorem 4.9 to see $d \geq |\mathcal{C}(y)|$. Since by the first part of the theorem $d \leq |\mathcal{C}(y)|$ we obtain $d = |\mathcal{C}(y)|$ when y is recurrent. $\square$

Corollary 4.23. *Let $\pi : X \rightarrow Y$ be a factor code from a SFT X to an irreducible sofic shift Y . Then*

- (a) $c_\pi = c_\pi^*$.
- (b) *There are exactly c_π transition classes over every right transitive point of Y .*
- (c) *Given an ergodic measure ν on Y , ν -almost every point of y has the same number of transition classes over it. We call this number $|\mathcal{C}(\nu)|$.*

Proof. Let (W, n, M) be a transition block of π of depth c_π^* and let y be a right transitive point of Y . Notice that W occurs infinitely often in the positive coordinates of y . Then (a) and (b) are clear by Theorem 4.22.

Now let ν be an ergodic measure on Y . Let (W', n', M') be a transition block of π with $\nu[W'] > 0$ such that

$$|M'| = \min\{|\tilde{M}| : (\tilde{W}, \tilde{n}, \tilde{M}) \text{ is a transition block of } \pi, \nu[\tilde{W}] > 0\}.$$

Note that W' occurs infinitely often in the positive coordinates of ν -almost every point of Y . Thus (c) is obtained as a direct corollary of Theorem 4.22. $\square$

We now show that for a finite-to-one factor code, the degree and the class degree of the code are the same.

Theorem 4.24. *Let $\pi : X \rightarrow Y$ be a finite-to-one factor code from a SFT to an irreducible sofic shift Y . Then $c_\pi = d_\pi$.*

Proof. Corollary 4.2 and Proposition 4.8(b) imply that the degree of a code and the class degree of a code are both invariant under conjugacy of factor triples. So by recoding, without loss of generality, we may assume X is a 1-step SFT and π is a 1-block factor code. Let y be a transitive point of Y . By Proposition 4.4 the preimages of y are mutually separated. It follows that each transition class above y has only one point. Therefore, $c_\pi = |\pi^{-1}(y)| = d_\pi$. $\square$

5 Bounding the number of ergodic measures of relative maximal entropy

In Section 3 we mentioned that although every 1-dimensional irreducible SFT has a unique ergodic measure (Parry measure) of maximal entropy, there can be more than one ergodic measure of relative maximal entropy over an ergodic measure; i.e., given a factor triple (X, Y, π) and an ergodic measure ν on Y , there can exist more than one ergodic measure on X that projects to ν under π and has maximal entropy among measures in the fibre $\pi^{-1}\{\nu\}$, see [17, Example 3.3]. In this section we show that the number of such measures over a fully supported ergodic measure can be no more than the class degree.

Theorem 5.1. *Let (X, Y, π) be a factor triple and ν be a fully supported ergodic measure on Y . The number of ergodic measures of relative maximal entropy over ν is at most c_π .*

Since entropy is a conjugacy invariant the following observation follows immediately.

Observation 5.2. *Let (X, Y, π) and $(\tilde{X}, \tilde{Y}, \tilde{\pi})$ be conjugate factor triples. Let ν be an ergodic measure on Y and $\tilde{\nu}$ be its corresponding ergodic measure on $\tilde{Y}$. The number of ergodic measures of relative maximal entropy over ν is the same as the number of ergodic measures of relative maximal entropy over $\tilde{\nu}$.*

In 2003, Petersen, Quas, and Shin [17] found an upper bound on the number of measures of relative maximal entropy. Let $N_\nu(\pi)$ denote the minimum number of symbols in $\pi_b^{-1}(w)$ as w runs over the symbols in $\mathcal{A}(Y)$ for which $\nu[w] > 0$.

Theorem 5.3. *[17, Corollary 1] Let $\pi : X \rightarrow Y$ be a 1-block code from a 1-step SFT X to a sofic shift Y . Let ν be an ergodic measure on Y . The number of ergodic measures of maximal entropy over ν is at most $N_\nu(\pi)$.*

This bound suffers from not being invariant under conjugacy. For example, the full 2-shift and any higher block presentation of it give different bounds on the number of ergodic measures of maximal entropy which map to the trivial measure on the full 1-shift.

One possibility to avoid having a non-invariant upper bound is to take the minimum of the bound in [17] over all conjugate factor triples with 1-block factor codes. This even improves the original bound. However, there is no known algorithm for computing this minimum.

Here we show that given an ergodic measure ν on Y , the number of transition classes over a ν -generic point of Y is an upper bound on the number of measures of relative maximal entropy over ν . Proposition 4.8(b) verifies that this bound is invariant under conjugacy of factor triples and Proposition 5.4 shows that it beats the bound mentioned above obtained by minimizing the bound in [17] over conjugate factor triples.

Proposition 5.4. *Let $\pi : X \rightarrow Y$ be a 1-block factor code from a SFT X to a sofic shift Y . Let ν be an ergodic measure on Y . Then we have*

$|\mathcal{C}_\pi(y)| \leq \min\{N_\nu(\tilde{\pi}) : (X, Y, \pi) \cong (\tilde{X}, \tilde{Y}, \tilde{\pi}), \tilde{\pi} \text{ is 1-block}\}$ where y is a ν -generic point of Y . Equality holds if π is finite-to-one and ν is fully supported.

Proof. Let the minimum of $N_\nu(\tilde{\pi})$ over all factor triples $(\tilde{X}, \tilde{Y}, \tilde{\pi})$ which are conjugate to (X, Y, π) and $\tilde{\pi}$ is 1-block, be attained at $(\bar{X}, \bar{Y}, \bar{\pi})$ where X is conjugate to $\bar{X}$ under a conjugacy $\phi : X \rightarrow \bar{X}$, Y is conjugate to $\bar{Y}$ under a conjugacy $\psi : Y \rightarrow \bar{Y}$, and $\bar{\pi} \circ \phi = \psi \circ \pi$. Denote the measure $\nu \circ \psi^{-1}$ by $\bar{\nu}$. Suppose that the minimum of $|\bar{\pi}_b^{-1}(w)|$ over all w in $\mathcal{A}(\bar{Y})$ with $\bar{\nu}[w] > 0$ is attained at a symbol $\bar{w}$. Let $\bar{y}$ be a $\bar{\nu}$ -generic point of $\bar{Y}$. There is a strictly increasing sequence of integers $(a_i)_{i \in \mathbb{N}}$ with $\bar{y}_{a_i} = \bar{w}$. Theorem 4.10 implies that $|\mathcal{C}_{\bar{\pi}}(\bar{y})| \leq |\bar{\pi}_b^{-1}(\bar{w})|$. Let $y = \psi^{-1}(\bar{y})$. Then by Proposition 4.8(a) we have $|\mathcal{C}_\pi(y)| = |\mathcal{C}_{\bar{\pi}}(\bar{y})| \leq |\bar{\pi}_b^{-1}(\bar{w})|$ which completes the proof of the first part of the result.

Now suppose π is finite-to-one and ν is fully supported. Since ν is fully supported we have

$$N_\nu(\tilde{\pi}) = \min\{|\tilde{\pi}_b^{-1}(w)| : w \in \mathcal{A}(\tilde{Y})\}.$$

Therefore the minimum of $N_\nu(\tilde{\pi})$ over all $(\tilde{X}, \tilde{Y}, \tilde{\pi})$ which are conjugate to (X, Y, π) and $\tilde{\pi}$ is 1-block is attained at a factor triple $(\bar{X}, \bar{Y}, \bar{\pi})$ with a magic symbol $\bar{w} \in \mathcal{A}(\bar{Y})$. By Proposition 4.4 we have $|\bar{\pi}_b^{-1}(\bar{w})| = d_{\bar{\pi}}$ which implies

$$\min\{N_\nu(\tilde{\pi}) : (X, Y, \pi) \cong (\bar{X}, \bar{Y}, \bar{\pi}), \bar{\pi} \text{ is 1-block}\} = d_{\bar{\pi}}.$$

Since the degree of a code is invariant under conjugacy of factor triples we deduce that

$$\min\{N_\nu(\tilde{\pi}) : (X, Y, \pi) \cong (\bar{X}, \bar{Y}, \bar{\pi}), \bar{\pi} \text{ is 1-block}\} = d_\pi.$$

Recalling that when π is finite-to-one the degree and the class degree of π are equal, the proof is complete. $\square$

Note that equality in Proposition 5.4 does not always hold. For example, consider the trivial factor code $\pi : \{0, 1\}^{\mathbb{Z}} \rightarrow \{0\}^{\mathbb{Z}}$. Then $c_\pi = 1$; however, if $(\tilde{X}, \{0\}^{\mathbb{Z}}, \tilde{\pi})$ is a factor triple conjugate to $(\{0, 1\}^{\mathbb{Z}}, \{0\}^{\mathbb{Z}}, \pi)$ then $\mathcal{A}(\tilde{X})$ must be strictly greater than 1 and therefore $N_\nu(\tilde{\pi}) > 1$ for any ergodic measure ν on Y .

Definition 5.5. Let $\pi : X \rightarrow Y$ be a 1-block factor code from a SFT X to a sofic shift Y and ν be an ergodic measure on Y . Let $\mu_1, \dots, \mu_n$ be invariant measures in the fibre $\pi^{-1}\{\nu\}$. The *relatively independent joining* $\tilde{\mu} = \mu_1 \otimes \dots \otimes \mu_n$ of $\mu_1, \dots, \mu_n$ over ν is defined as follows: if $A_1, \dots, A_n$ are measurable subsets of X then

$$\hat{\mu}(A_1 \times \dots \times A_n) = \int_Y \prod_{i=1}^n \mathbb{E}_{\mu_i}(\mathbf{1}_{A_i} | \pi^{-1} \mathcal{B}_Y) \circ \pi^{-1} d\nu.$$

Writing p_i for the projection $X^n \rightarrow X$ onto the i th coordinate, it is shown that for $\hat{\mu}$ -almost every $\hat{x} \in X^n$, $\pi(p_i(\hat{x}))$ is the same for each i [18].

We will use Theorem 5.6 below which is the main theorem from [17] to prove a stronger theorem (Theorem 5.7).

Theorem 5.6. [17, Theorem 1] Let $\pi : X \rightarrow Y$ be a 1-block factor code from a 1-step SFT X to a sofic shift Y . Let ν be an ergodic measure on Y , and two distinct ergodic measures μ_1 and μ_2 be measures of relative maximal entropy over ν . Then $(\mu_1 \otimes \mu_2)\{(u, v) \in X \times X : u_0 = v_0\} = 0$.

Theorem 5.7. Let (X, Y, π) be a factor triple. Let ν be an ergodic measure on Y , and two distinct measures μ_1 and μ_2 be ergodic measures of relative maximal entropy over ν . Then $(\mu_1 \otimes \mu_2)\{(u, v) \in X \times X : u \sim v\} = 0$.

Proof. First we show that, without loss of generality, we may assume X is a 1-step SFT and π is a 1-block factor code with a magic symbol. Suppose $(\tilde{X}, \tilde{Y}, \tilde{\pi})$ is a factor triple conjugate to (X, Y, π) and $\phi : X \rightarrow \tilde{X}$ is a conjugacy from X to $\tilde{X}$. By the proof of Proposition 4.8(a) we have

$$(\phi \times \phi)\{(u, v) \in X \times X : u \sim v\} = \{(\phi(u), \phi(v)) \in \tilde{X} \times \tilde{X} : \phi(u) \sim \phi(v)\}.$$

Moreover, the corresponding measure to $\mu_1 \otimes \mu_2$ under the conjugacy

$$\phi \times \phi : X \times X \rightarrow \tilde{X} \times \tilde{X},$$

is $\tilde{\mu}_1 \otimes \tilde{\mu}_2$ where $\tilde{\mu}_1$ and $\tilde{\mu}_2$ are corresponding measures to μ_1 and μ_2 under ϕ . It follows that

$$(\mu_1 \otimes \mu_2)\{(u, v) \in X \times X : u \sim v\} = (\tilde{\mu}_1 \otimes \tilde{\mu}_2)\{(\phi(u), \phi(v)) \in \tilde{X} \times \tilde{X} : \phi(u) \sim \phi(v)\}.$$

Therefore by recoding, without loss of generality, we may assume X is a 1-step SFT and π is a 1-block factor code.

Let u, v be in X , n be in $\mathbb{Z}$, and a be a symbol in $\mathcal{A}(X)$. Write $u \overset{a}{\approx}_n v$ if $u \sim v$ and there are integers $m \leq n$ and $p \geq n$ such that

$$v_i = \begin{cases} u_i & -\infty < i < m, p < i < \infty \\ a & i = n \end{cases}$$

It is worth mentioning that the relation $\overset{a}{\approx}_n$ is not in general an equivalence relation on X . We have

$$\{(u, v) : u \sim v\} = \bigcup_{a \in \mathcal{A}(X)} \bigcup_{n \in \mathbb{Z}} \{(u, v) : u \overset{a}{\approx}_n v\}.$$

Now suppose, for a contradiction, that $(\mu_1 \otimes \mu_2)\{(u, v) : u \sim v\} > 0$. There must be an integer n and a symbol a in $\mathcal{A}(X)$ such that

$$(\mu_1 \otimes \mu_2)\{(u, v) : u \overset{a}{\approx}_n v\} > 0.$$

By applying $(T \times T)^{-n}$, without loss of generality, we may assume $n = 0$. Since we have

$$\{(u, v) : u \overset{a}{\approx}_0 v\} = \bigcup_{k \in \mathbb{Z}} \{(u, v) : u \overset{a}{\approx}_0 v, u_{-k}^k, v_{-k}^k \text{ are routable through } a \text{ at time } 0\},$$

it follows that there is an integer k so that

$$(\mu_1 \otimes \mu_2)\{(u, v) : u \overset{a}{\approx}_0 v, u_{-k}^k, v_{-k}^k \text{ are routable through } a \text{ at time } 0\} > 0.$$

Considering only blocks of length $2k + 1$, there must be blocks A, B in X such that

$$(\mu_1 \otimes \mu_2)\{(u, v) : u \overset{a}{\approx}_0 v, u_{[-k, k]} = A, v_{[-k, k]} = B\} > 0. \quad (5.1)$$

Both A and B are routable through a at time n ; i.e., there are blocks A', B' in $\pi_b^{-1}(W)$ with $A'_{-k} = A_{-k}$, $B'_{-k} = B_{-k}$, $A'_0 = B'_0 = a$, $A'_k = A_k$, and $B'_k = B_k$. Let $G \subseteq \mathbb{Z}$ be the set $\{-k, \dots, k\}$. Basic properties of conditional expectation imply that

$$\begin{aligned} \mathbb{E}_{\mu_1}(\mathbf{1}_{[A]} | \pi^{-1}(\mathcal{B}_Y)) &= \mathbb{E}(\mathbb{E}_{\mu_1}(\mathbf{1}_{-n[A]} | \pi^{-1}(\mathcal{B}_Y) \vee \sigma(X_{G^c})) | \pi^{-1}(\mathcal{B}_Y)) \\ &= \mathbb{E}(\mathbb{E}_{\mu_1}(\mathbf{1}_{-n[A']} | \pi^{-1}(\mathcal{B}_Y) \vee \sigma(X_{G^c})) | \pi^{-1}(\mathcal{B}_Y)) \quad (5.2) \\ &= \mathbb{E}_{\mu_1}(\mathbf{1}_{[A']} | \pi^{-1}(\mathcal{B}_Y)), \end{aligned}$$

where the second equality follows from Theorem 3.3. Similarly we have

$$\mathbb{E}_{\mu_2}(\mathbf{1}_{[B]}|\pi^{-1}\mathcal{B}_Y) = \mathbb{E}_{\mu_2}(\mathbf{1}_{[B']}|\pi^{-1}\mathcal{B}_Y). \quad (5.3)$$

Let $D = \{(u, v) : u \stackrel{a}{\approx}_0 v, u_{[-k, k]} = A, v_{[k, k]} = B\}$. Since $D \subseteq {}_{-k}[A] \times {}_{-k}[B]$ we have

$$\begin{aligned} \mu(D) &\leq (\mu_1 \otimes \mu_2)([A] \times [B]) \\ &= \int_Y \mathbb{E}_{\mu_1}(\mathbf{1}_{[A]}|\pi^{-1}\mathcal{B}_Y) \mathbb{E}_{\mu_2}(\mathbf{1}_{[B]}|\pi^{-1}\mathcal{B}_Y) \circ \pi^{-1} d\nu \\ &= \int_Y \mathbb{E}_{\mu_1}(\mathbf{1}_{[A']}|\pi^{-1}\mathcal{B}_Y) \mathbb{E}_{\mu_2}(\mathbf{1}_{[B']}|\pi^{-1}\mathcal{B}_Y) \circ \pi^{-1} d\nu \quad \text{using (5.2) and (5.3)} \\ &= (\mu_1 \otimes \mu_2)([A'] \times [B']) \\ &= 0 \end{aligned}$$

where the last equality follows from Theorem 5.6 since $A'_0 = B'_0 = a$. This contradicts Equation (5.1). $\square$

The following theorem is a more general case of Theorem 5.1.

Theorem 5.8. *Let (X, Y, π) be a factor triple and ν be an ergodic measure on Y . The number of ergodic measures of relative maximal entropy over ν is at most $|\mathcal{C}(\nu)|$.*

Proof. Let $(\tilde{X}, \tilde{Y}, \pi)$ be a factor triple conjugate to (X, Y, π) under conjugacies $\phi : X \rightarrow \tilde{X}$ and $\psi : Y \rightarrow \tilde{Y}$. By Observation 5.2, the number of ergodic measures of relative maximal entropy over ν and $\nu \circ \psi^{-1}$ is the same. Moreover, by Proposition 4.8(a), for each y in Y we have $|\mathcal{C}_\pi(y)| = |\mathcal{C}_{\tilde{\pi}}(\psi(y))|$. Therefore by recoding, without loss of generality, we may assume X is a 1-step SFT and π is a 1-block factor code.

Suppose, for a contradiction, that there are $n > |\mathcal{C}(\nu)|$ ergodic measures $\mu_1, \dots, \mu_n$ on X of relative maximal entropy over ν . Form the relatively independent joining $\hat{\mu}$ on X^n of the measures $\mu_1, \dots, \mu_n$. Note that for $\hat{\mu}$ -a.e. $\hat{x} \in X^n$, $\pi(p_i(\hat{x}))$ has $|\mathcal{C}(\nu)|$ transition classes over it. The assumption $n > |\mathcal{C}(\nu)|$ implies that for $\hat{\mu}$ -a.e. $\hat{x} = (x_1, \dots, x_n) \in X^n$ there are distinct i, j such that $p_i(\hat{x}) \sim p_j(\hat{x})$; i.e.,

$$\hat{\mu} \left(\bigcup_{1 \leq i < j \leq n} \{\hat{x} = (x_1, \dots, x_n) : x_i \sim x_j\} \right) = 1.$$

At least one of the sets $S_{i,j} = \{(x_1, \dots, x_n) : x_i \sim x_j\}$ must have positive $\hat{\mu}$ -measure. It follows that

$$\begin{aligned} 0 &< \hat{\mu}(S_{i,j}) \\ &= \hat{\mu}(\{(x_1, \dots, x_n), x_i \sim x_j\}) \\ &= (\mu_i \otimes \mu_j)\{(u, v) : u \sim v\}. \end{aligned}$$

This contradicts Theorem 5.7. □

6 Open Questions

Let X be a one-sided topologically mixing shift of finite type. An invariant measure μ on X is a **Gibbs measure** corresponding to $f \in C(X)$ if there are constants $C_1, C_2 > 0$ and $P > 0$ such that

$$C_1 \leq \frac{\mu([x_0 x_1 \dots x_{n-1}])}{\exp(-Pn + (S_n f)(x))} \leq C_2$$

for every $x \in X$ and $n \geq 1$, where $(S_n f)(x) = \sum_{k=0}^{n-1} f(T^k(x))$.

Walters [24, 25] introduces a class $\text{Bow}(X)$ of functions that contains the functions with summable variation, all of which have unique equilibrium states. Let

$$\text{var}_n(f) = \sup\{|f(x) - f(y)| : x, y \in X, x_i = y_i \text{ for all } 0 \leq i \leq n-1\}.$$

Then $\text{Bow}(X, T) = \{f \in C(X) : \sup_{n \geq 1} \text{var}_n(S_n f) < \infty\}$.

Theorem 6.1. [24, Theorem 2.16] *Let $f \in \text{Bow}(X)$. Then f has a unique equilibrium state μ which is a Gibbs measure.*

The relative version of this result is an open question. We make the following conjecture.

Conjecture 6.2. *Let (X, Y, π) be a factor triple and ν be a fully supported ergodic measure on Y . For any function $f \in \text{Bow}(X)$ the number of ergodic measures of maximal pressure of f in the fibre $\pi^{-1}\{\nu\}$ is at most c_π .*

References

- [1] D. Blackwell. The entropy of functions of finite-state Markov chains. In *Transactions of the first Prague conference on information theory, Statistical decision functions, random processes held at Liblice near Prague from November 28 to 30, 1956*, pages 13–20. Publishing House of the Czechoslovak Academy of Sciences, Prague, 1957.
- [2] M. Boyle. Putnam’s resolving maps in dimension zero. *Ergodic Theory Dynam. Systems*, 25(5):1485–1502, 2005. See erratum at <http://www-users.math.umd.edu/~mmb/papers/degreeerratum>.
- [3] M. Boyle and S. Tuncel. Infinite-to-one codes and Markov measures. *Trans. Amer. Math. Soc.*, 285(2):657–684, 1984.
- [4] C. J. Burke and M. Rosenblatt. A Markovian function of a Markov chain. *Ann. Math. Statist.*, 29:1112–1122, 1958.
- [5] R. Burton and J. E. Steif. Non-uniqueness of measures of maximal entropy for subshifts of finite type. *Ergodic Theory Dynam. Systems*, 14(2):213–235, 1994.
- [6] D. Gatzouras and Y. Peres. The variational principle for Hausdorff dimension: a survey. In *Ergodic theory of $\mathbf{Z}^d$ actions (Warwick, 1993–1994)*, volume 228 of *London Math. Soc. Lecture Note Ser.*, pages 113–125. Cambridge Univ. Press, Cambridge, 1996.
- [7] D. Gatzouras and Y. Peres. Invariant measures of full dimension for some expanding maps. *Ergodic Theory Dynam. Systems*, 17(1):147–167, 1997.
- [8] R. B. Israel. *Convexity in the theory of lattice gases*. Princeton University Press, Princeton, N.J., 1979. Princeton Series in Physics, With an introduction by Arthur S. Wightman.
- [9] G. Keller. *Equilibrium states in ergodic theory*, volume 42 of *London Mathematical Society Student Texts*. Cambridge University Press, Cambridge, 1998.

- [10] O. E. Lanford, III and D. Ruelle. Observables at infinity and states with short range correlations in statistical mechanics. *Comm. Math. Phys.*, 13:194–215, 1969.
- [11] F. Ledrappier and P. Walters. A relativised variational principle for continuous transformations. *J. London Math. Soc. (2)*, 16(3):568–576, 1977.
- [12] F. Ledrappier and L.-S. Young. The metric entropy of diffeomorphisms. II. Relations between entropy, exponents and dimension. *Ann. of Math. (2)*, 122(3):540–574, 1985.
- [13] D. Lind and B. Marcus. *An introduction to symbolic dynamics and coding*. Cambridge University Press, Cambridge, 1995.
- [14] B. Marcus, K. Petersen, and S. Williams. Transmission rates and factors of Markov chains. In *Conference in modern analysis and probability (New Haven, Conn., 1982)*, volume 26 of *Contemp. Math.*, pages 279–293. Amer. Math. Soc., Providence, RI, 1984.
- [15] W. Parry. Intrinsic Markov chains. *Trans. Amer. Math. Soc.*, 112:55–66, 1964.
- [16] K. Petersen. Information compression and retention in dynamical processes. In *Dynamics and randomness (Santiago, 2000)*, volume 7 of *Non-linear Phenom. Complex Systems*, pages 147–217. Kluwer Acad. Publ., Dordrecht, 2002.
- [17] K. Petersen, A. Quas, and S. Shin. Measures of maximal relative entropy. *Ergodic Theory Dynam. Systems*, 23(1):207–223, 2003.
- [18] D. J. Rudolph. *Fundamentals of measurable dynamics*. Oxford Science Publications. The Clarendon Press Oxford University Press, New York, 1990. Ergodic theory on Lebesgue spaces.
- [19] D. Ruelle. Statistical mechanics on a compact set with Z^v action satisfying expansiveness and specification. *Trans. Amer. Math. Soc.*, 187:237–251, 1973.
- [20] C. E. Shannon. A mathematical theory of communication. *Bell System Tech. J.*, 27:379–423, 623–656, 1948.

- [21] S. Shin. Measures that maximize weighted entropy for factor maps between subshifts of finite type. *Ergodic Theory Dynam. Systems*, 21(4):1249–1272, 2001.
- [22] P. Walters. *An introduction to ergodic theory*, volume 79 of *Graduate Texts in Mathematics*. Springer-Verlag, New York, 1982.
- [23] P. Walters. Relative pressure, relative equilibrium states, compensation functions and many-to-one codes between subshifts. *Trans. Amer. Math. Soc.*, 296(1):1–31, 1986.
- [24] P. Walters. Convergence of the Ruelle operator for a function satisfying Bowen’s condition. *Trans. Amer. Math. Soc.*, 353(1):327–347 (electronic), 2001.
- [25] P. Walters. Regularity conditions and Bernoulli properties of equilibrium states and g -measures. *J. London Math. Soc. (2)*, 71(2):379–396, 2005.

